
## **Supplementary information:**

Bone metastasis is associated with acquisition of mesenchymal phenotype and immune suppression in a model of spontaneous breast cancer metastasis

Lea Monteran<sup>1</sup>, Nour Ershaid<sup>1</sup>, Idan Sabah<sup>1</sup>, Ibrahim Fahoum<sup>2</sup>, Yael Zait<sup>1</sup>, Ophir Shani<sup>1</sup>, Noam Cohen<sup>1</sup>, Anat Eldar-Boock<sup>3</sup>, Ronit Satchi-Fainaro<sup>3</sup> and Neta Erez<sup>1\*</sup>

<sup>1</sup>Department of Pathology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv 69978, Israel <sup>2</sup>Department of Pathology, Tel Aviv Sourasky Medical Center, Tel Aviv University, Tel Aviv, Israel. <sup>3</sup>Department of Physiology and Pharmacology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv 69978, Israel

\*Correspondence to netaerez@tauex.tau.ac.il



Supplementary Figure 1: Breast cancer metastasis is associated with changes in the immune milieu in lungs. (A) Scheme of experimental design. This figure was designed by using graphical elements from BioRender. (B-M) FACS analysis quantification of major immune cell populations in the lungs of normal, 4T1 and 4T-Bone-injected mice. (B) CD45+CD11b+CD11c- total myeloid cell population. (C) CD45+CD11b+CD11c+Ly6Cint myeloid-dendritic cells. (D) CD45+CD11c+CD11b- dendritic cells. (E) CD45+CD11c-CD11b+Ly6G+Ly6Cint granulocytes. (F) CD45+B220-CD3-Nkp46+NK cells. (G) CD45+B220-CD3+ T cells. (H) CD45+CD11c+CD11b-SiglecF+Ly6C+ Alveolar Macrophages (AMs). (I) CD45+B220+CD3- B cells. (J) CD45+CD11c+CD11b+Ly6C- interstitial macrophages (IMs). (K) CD45+CD11c-CD11b+Ly6G-Ly6C+ monocytes. (L) CD45+CD11b+CD11cintLy6C-Ly6C- monocytes. (M) CD45+CD11c-CD11b+SiglecF+Ly6G- eosinophils. n=6-10 mice per group. Data are all presented as % of CD45, normalized to the average of normal group. Error bars represent SEM. p<0.05; Kruskal-Wallis Multiple comparison test.